

## ImPACT Reference List

### 2014

1 year test-retest reliability of ImPACT in professional ice hockey players. Bruce J, Echemendia R, Meeuwisse W, Comper P, Sisco A. *Clin Neuropsychol*. 2014;28(1):14-25.

Age and test setting affect the prevalence of invalid baseline scores on neurocognitive tests. Lichtenstein JD, Moser RS, Schatz P. *Am J Sports Med*. 2014 Feb;42(2):479-84.

Baseline evaluation in youth ice hockey players: comparing methods for documenting prior concussions and attention or learning disorders. McKay CD, Schneider KJ, Brooks BL, Mrazik M, Emery CA *J Orthop Sports Phys Ther*. 2014 May;44(5):329-35.

Brain activation during neurocognitive testing using functional near-infrared spectroscopy in patients following concussion compared to healthy controls. Kontos AP, Huppert TJ, Beluk NH, Elbin RJ, Henry LC, French J, Dakan SM, Collins MW. *Brain Imaging Behav*. 2014 Jan 30.

A brief vestibular/ocular motor screening (VOMS) assessment to evaluate concussions: preliminary findings. Mucha A, Collins MW, Elbin RJ, Furman JM, Troutman-Enseki C, DeWolf RM, Marchetti G, Kontos AP. *Am J Sports Med*. 2014 Aug 8.

C-41 changes in symptoms in concussed and non-concussed athletes following neuropsychological assessment. Meyer J, Arnett P. *Arch Clin Neuropsychol*. 2014 Sep;29(6):589.

C-45 Do athletes with dyslexia differ at baseline and/or at concussion post injury assessment on a computer-based test battery? Johnson E, Pardini J, Sandel N, Lovell M. *Arch Clin Neuropsychol*. 2014 Sep;29(6):590-1.

C-46A new iPad version of ImPACT is resistant to "sandbagging." Ferris C,

Schatz P. *Arch Clin Neuropsychol*. 2014 Sep;29(6):591.

Cerebral concussion: a historical perspective. Maroon JC, Mathyssek C, Bost J. *Prog Neurol Surg*. 2014;28:1-13.

Comparing post-concussive neurocognitive test data to normative data presents risks for under-classifying "above average" athletes. Schatz P, Robertshaw S. *Arch Clin Neuropsychol*. 2014 Sep 1. [Epub ahead of print].

Computerized neurocognitive testing within 1 week of sport-related concussion: meta-analytic review and analysis of moderating factors. Kontos AP, Braithwaite R, Dakan S, Elbin RJ. *J Int Neuropsychol Soc*. 2014 Mar;20(3):324-32.

Does mechanism of injury play a role in recovery from concussion? Seiger A, Goldwater E, Deibert E. *J Head Trauma Rehabil*. 2014 Jun 4.

Effect of sex on symptoms and return to baseline in sport-related concussion. Zuckerman SL, Apple RP, Odom MJ, Lee YM, Solomon GS, Sills AK. *J Neurosurg Pediatr*. 2014 Jan;13(1):72-81.

Examination of the test-retest reliability of a computerized neurocognitive test battery. Nakayama Y, Covassin T, Schatz P, Nogle S, Kovan J. *Am J Sports Med*. 2014 Jun 6;42(8):2000-2005.

Examining postconcussion symptoms of dizziness and imbalance on neurocognitive performance in collegiate football players. Honaker JA, Lester HF, Patterson JN, Jones SM. *Otol Neurotol*. 2014 Jul;35(6):1111-7.

Neurocognitive performance and symptom profiles of Spanish-speaking Hispanic athletes on the ImPACT test. Ott S, Schatz P, Solomon G, Ryan JJ. *Arch Clin Neuropsychol*. 2014 Mar;29(2):152-63.

Neuropsychological factors related to college ice hockey concussions. Pedersen HA, Ferraro FR, Himle M, Schultz C, Poolman M. *Am J Alzheimers Dis Other Demen*. 2014 May;29(3):201-4.

Neuropsychological test performance of Hawai'i high school athletes:

updated Hawai'i immediate post-concussion assessment and cognitive testing data. Tsushima WT, Siu AM. *Hawaii J Med Public Health*. 2014 Jul;73(7):208-11.

A potential biomarker in sports-related concussion: Brain functional connectivity alteration of the default-mode network measured with longitudinal resting-state fMRI over 30 days. Zhu DC, Covassin T, Nogle S, Doyle S, Russell D, Pearson RL, Monroe J, Liszewski CM, DeMarco JK, Kaufman DI. *J Neurotrauma*. 2014 Aug 12.

Psychometric properties and reference values for the ImPACT neurocognitive test battery in a sample of elite youth ice hockey players. McKay CD, Brooks BL, Mrazik M, Jubinville AL, Emery CA. *Arch Clin Neuropsychol*. 2014 Mar;29(2):141-51.

Relationship between concussion history and neurocognitive test performance in National Football League draft picks. Solomon GS, Kuhn A. *Am J Sports Med*. 2014 Apr;42(4):934-9.

A review of modifying factors affecting usage of diagnostic rating scales in concussion management. Dessy A, Rasouli J, Gometz A, Choudhri T. *Clin Neurol Neurosurg*. 2014 Jul;122:59-63.

Screening for lifetime concussion in athletes: importance of oculomotor measures. Poltavski DV, Biberdorf D. *Brain Inj*. 2014;28(4):475-85.

Sideline assessment tools for the evaluation of concussion in athletes: A review. Okonkwo D, Tempel Z, Maroon J. *J Neuropsychology*. 2014 Oct.

The underreporting of self-reported symptoms following sports-related concussion. Meier TB, Brummel BJ, Singh R, Nerio CJ, Polanski DW, Bellgowan PS. *J Sci Med Sport*. 2014 Jul 24. pii: S1440-2440(14)00134-0.

Utility of repeated assessment after invalid baseline performance. Schatz P, Kelley T, Ott SD, Solomon GS, Elbin RJ, Higgins K, Moser RS. *J Athl Train*. 2014 Aug 27.

## 2013

- AMPA peptide values in blood of non-athletes and club sport athletes with concussions. Dambinova SA, Shikuev, Weissman JD, Mullins, JD. *Military Medicine*. 2013, 178 (3):285-290.
- Baseline neurocognitive scores in athletes with attention deficit-spectrum disorders and/or learning disability. Zuckerman SL, Lee YM, Odom MJ, Solomon GS, Sills AK. *J Neurosurg Pediatr*. 2013.
- Computerized assessment of cognitive late effects among adolescent brain tumor survivors. Conklin HM, Ashford JM, Di Pinto M, Vaughn CG, Merchant TW, Ogg RJ, Santana V, Wu, S. *J Neurooncol*. 2013 (Epub ahead of print).
- Concussion symptoms and neurocognitive performance of high school and college athletes who incur multiple concussions. Covassin T, Moran R, Wilhelm K. *Am J Sports Med*. 2013.
- Discriminant construct validity of ImPACT: A companion study. Maerlender A, Flashman L, Kessler A, et al. *Clin Neuropsychol*. 2013;27(2):290-299.
- Effects of youth football on selected clinical measures of neurologic function: A pilot study. Munce TA, Dorman JC, Odney TO, Thompson PA, Valentine VD, Bergeron MF. *J Child Neurol*. 2013 Nov 21.
- Higher BMI is associated with reduced cognitive performance in division I athletes. Fedor A, Gunstad J. *Obes Facts*. 2013;6(2):185-192.
- ImPACT test-retest reliability: Reliably unreliable? Resch J, Driscoll A, McCaffrey N, et al. *J Athl Train*. 2013.
- Individual and combined effects of LD and ADHD on computerized neurocognitive concussion test performance: Evidence for separate norms. Elbin RJ, Kontos AP, Kegel N, Johnson E, Burkhart S, Schatz P. *Arch Clin Neuropsychol*. 2013.

- Intraindividual cognitive variability before and after sports-related concussion. Rabinowitz AR, Arnett PA. *Neuropsychology*. 2013;27(4):481-490.
- Management strategies and medication use for treating pediatric patients with concussions. Kinnaman KA, Mannix RC, Comstock RD, Meehan WP. *Acta Paediatr*. 2013.
- Neurocognitive functioning and symptoms reporting of high school athletes following a single concussion. Tsushima WT, Shirakawa N, Geling O. *Appl Neuropsychol Child*. 2013;2(1):13-16.
- Neurocognitive responses to a single session of static squats with whole body vibration. Amonette WE, Boyle M, Psarakis M, Barker J, Dupler TL, Ott S. *The Journal of Strength & Conditioning Research*. 2013.
- Neurocognitive test performance and symptom reporting in cheerleaders with concussions. Lovell MR, Solomon GS. *J Pediatr*. 2013.
- One-month test-retest reliability of the ImPACT Test battery Schatz P, Ferris CS. *Arch Clin Neuropsychol*. 2013 May 23. (Epub ahead of print).
- Post-exertion neurocognitive test failure among student-athletes following concussion. McGrath N, Dinn WM, Collins MW, Lovell MR, Elbin RJ, Kontos AP. *Brain Inj*. 2013;27(1):103-113.
- Posttraumatic migraine as a predictor of recovery and cognitive impairment after sport-related concussion Kontos AP, Elbin RJ, Lau B, Simensky S, Freund B, French J, Collins MW. *Am J Sports Med*. 2013 May 22. 2013 May 22. (Epub ahead of print).
- Practice effects reveal visuomotor vulnerability in school and university rugby players. Shuttleworth-Edwards AB, Radloff SE, Whitefield-Alexander VJ, Smith IP, Horsman M. *Arch Clin Neuropsychol*. 2013.
- The prevalence of undiagnosed concussions in athletes. Meehan WP,3rd, Mannix RC, O'brien MJ, Collins MW. *Clin J Sport Med*. 2013.
- A prospective study of symptoms and neurocognitive outcomes in youth

with concussion vs orthopaedic injuries. Rieger BP, Lewandowski LJ, Callahan JM, et al. *Brain Inj.* 2013;27(2):169-178.

The relationship among neurocognitive performances and biopsychosocial characteristics of elite national football league draft picks: An exploratory investigation. Solomon GS, Haase RF, Kuhn A. *Arch Clin Neuropsychol.* 2013;28(1):9-20.

The relationship of symptoms and neurocognitive performance to perceived recovery from sports-related concussion among adolescent athletes. Sandel NK, Lovell MR, Kegel NE, Collins, MW, Kontos AP. *Appl. Neuropsychol Child.* 2013;2(1):64-69.

“Sandbagging” baseline test performance on ImPACT without detection is more difficult than it appears. Schatz P, Glatts C. *Arch. Clin Neuropsychol.* 2013, Feb 11.

Sensitivity and specificity of the online version of ImPACT in high school and collegiate athletes. Schatz P, Sandel N. *Am. J. Sports Med.* 2013;41(2):321-326.

Sport concussion knowledge base, clinical practises and needs for continuing medical education: A survey of family physicians and cross-border comparison. Lebrun CM, Mrazik M, Prasad AS, et al. *Br J Sports Med.* 2013;47(1):54-59.

Subjective, but not objective, lingering effects of multiple past concussions in adolescents. Brooks B, McKay C, Mrazik M, Barlow K, Meeuwisse W, Emery, C. *J. Neurotrauma.* 2013 (Epub ahead of print).

Symptom severity predicts prolonged recovery after sport-related concussion, but age and amnesia do not. Meehan WP, 3rd, Mannix RC, Stracciolini A, Elbin RJ, Collins MW. *J Pediatr.* 2013.

Symptomatic white matter changes in mild traumatic brain injury resemble pathologic features of early alzheimer dementia. Fakhran S, Yaeger K, Alhilali L. *Radiology.* 2013.

Test-retest reliability of four computerized neurocognitive assessment tools

in an active duty military population. Cole WR, Arrieux JP, Schwab K, Ivins BJ, Qashu FM, Lewis SC. *Arch Clin Neuropsychol* 2013.

Use of ImPACT to diagnose minimal hepatic encephalopathy: An accurate, practical and user-friendly Internet-based test battery. Tsushima M, Tsushima WT, Tsushima V, Lim N, Madrigal E, Jackson C, Medler MH. *Dig Dis Sci*. 2013. (Epub ahead of print).

## 2012

145 sport-related concussion and age: Number of days to neurocognitive baseline. Zuckerman SL, Odom M, Lee YM, Forbes J, Sills AK, Solomon G. *Neurosurgery*. 2012;71(2):E558.

Age-related differences and reliability on computerized and paper-and-pencil neurocognitive assessment batteries. Register-Mihalik JK, Kontos DL, Guskiewicz KM, Mihalik JP, Conder R, Shields EW. *J Athl Train*. 2012;47(3):297-305.

Are there gender differences in cognitive function, chronic stress, and neurobehavioral symptoms after mild-to-moderate traumatic brain injury? Covassin T, Bay E. *J Neurosci Nurs*. 2012;44(3):124-133.

A case for mental and physical rest in youth sports concussion: it's never too late. Moser RS, Schatz P. *Front. Neurol*. 2012;3:171.

Clinical utility of ImPACT assessment for post-concussion return-to-play counseling: Psychometric issues. Mayers LB, Redick TS. *J Clin Exp Neuropsychol*. 2012;34(3):235-242.

Cognitive effects of one season of head impacts in a cohort of collegiate contact sport athletes. McAllister TW, Flashman LA, Maerlender A, et al. *Neurology*. 2012;78(22):1777-1784.

Computerized neurocognitive testing for the management of sport-related concussions. Meehan WP 3rd, d'Hemecourt P, Collins CL, Taylor AM, Comstock RD. *Pediatrics*. 2012;129(1):38-44.

Concussions: What a neurosurgeon should know about current scientific

evidence and management strategies. Neal MT, Wilson JL, Hsu W, Powers AK. *Surg Neurol Int.* 2012;3:16.

Cutoff scores in neurocognitive testing and symptom clusters that predict protracted recovery from concussions in high school athletes. Lau BC, Collins MW, Lovell MR. *Neurosurgery.* 2012;70(2):371-9; discussion 379.

Depression and neurocognitive performance after concussion among male and female high school and collegiate athletes. Kontos AP, Covassin T, Elbin RJ, Parker T. *Arch Phys Med Rehabil.* 2012;93(10):1751-1756.

Diagnostic efficiency of ImPACT and CogSport in concussed rugby union players who have not undergone baseline neurocognitive testing. Gardner A, Shores EA, Batchelor J, Honan CA. *Appl Neuropsychol.* 2012;19(2):90-97.

Do brain activation changes persist in athletes with a history of multiple concussions who are asymptomatic? Elbin RJ, Covassin T, Hakun J, et al. *Brain Inj.* 2012;26(10):1217-1225.

The effect of hyperbaric oxygen on symptoms following mild traumatic brain injury. Wolf EG, Cifu D, Baugh L, Carne W, Profenna L. *J Neurotrauma.* 2012

Efficacy of amantadine treatment on symptoms and neurocognitive performance among adolescents following sports-related concussion. Reddy CC, Collins M, Lovell M, Kontos AP. *J Head Trauma Rehabil.* 2012

Efficacy of immediate and delayed cognitive and physical rest for treatment of sports-related concussion. Moser RS, Glatts C, Schatz P. *J Pediatr.* 2012

Epidemiology, trends, assessment and management of sport-related concussion in united states high schools Guerriero RM, Proctor MR, Mannix R, Meehan WP,3rd. *Curr Opin Pediatr.* 2012.

How many HITS are too many?: The use of accelerometers to study sports-



related concussion. Deibert E, Kryscio R. *Neurology*. 2012;78(22):1712-1713.

Identification and validation of prognostic criteria for persistence of mild traumatic brain injury-related impairment in the pediatric patient. Wiebe DJ, Collins MW, Nance ML. *Pediatr Emerg Care*. 2012;28(6):498-502.

Neuropsychological and neurophysiological assessment of sport concussion in children, adolescents and adults. Baillargeon A, Lassonde M, Leclerc S, Ellemberg D. *Brain Injury*. 2012;26(3):211-220.

Neuropsychological testing for sports-related concussion: How athletes can sandbag their baseline testing without detection. Erdal K. *Arch Clin Neuropsychol*. 2012;27(5):473-479.

Pediatric sports-related concussion produces cerebral blood flow alterations. Maugans TA, Farley C, Altaye M, Leach J, Cecil KM. *Pediatrics*. 2012;129(1):28-37.

Possible lingering effects of multiple past concussions. Iverson GL, Echemendia RJ, Lamarre AK, Brooks BL, Gaetz MB. *Rehabil Res Pract*. 2012;2012:316575.

Predictors of postconcussive symptoms 3 months after mild traumatic brain injury. Ponsford J, Cameron P, Fitzgerald M, Grant M, Mikocka-Walus A, Schonberger M. *Neuropsychology*. 2012;26(3):304-313.

Prevalence of invalid computerized baseline neurocognitive test results in high school and collegiate athletes. Schatz P, Moser RS, Solomon GS, Ott SD, Karpf R. *J Athl Train*. 2012;47(3):289-296.

Prevalence of neurocognitive and balance deficits in collegiate aged football players without clinically diagnosed concussion. Mulligan I, Boland M, Payette J. *J Orthop Sports Phys Ther*. 2012;42(7):625-632.

A prospective study of physician-observed concussion during a varsity university ice hockey season: Incidence and neuropsychological changes. part 2 of 4. Echlin PS, Skopelja EN, Worsley R, et al. *Neurosurg Focus*. 2012;33(6):E2: 1-11.

- Psychometric properties of self-report concussion scales and checklists. McLeod TC, Leach C. *J Athl Train*. 2012;47(2):221-223.
- Recovery from sports-related concussion: Days to return to neurocognitive baseline in adolescents versus young adults. Zuckerman SL, Lee YM, Odom MJ, Solomon GS, Forbes JA, Sills AK. *Surg Neurol Int*. 2012;3:130-7806.102945. Epub 2012 Oct 27.
- Residual effects of combat-related mild traumatic brain injury. Kontos AP, Kotwal RS, Elbin R, Lutz RH, Forsten RD, Benson PJ. *J Neurotrauma*. 2012
- Response to acute concussive injury in soccer players: Is gender a modifying factor? Zuckerman SL, Solomon GS, Forbes JA, Haase RF, Sills AK, Lovell MR. *J Neurosurg Pediatr*. 2012
- Response to Mayers and Redick: "Clinical utility of ImPACT assessment for postconcussion return-to-play counseling: Psychometric issues." Schatz P, Kontos A, Elbin R. *J Clin Exp Neuropsychol*. 2012;34(4):428-34; discussion 435-442.
- A revised factor structure for the post-concussion symptom scale: baseline and post-concussion factors. Kontos AP, Elbin RJ, Schatz P, Covassin T, Henry L, Pardini, J, Collins, MW. *Am. J Sports Med*. 2012 Oct 40(10):2375-2384.
- The role of age and sex in symptoms, neurocognitive performance, and postural stability in athletes after concussion. Covassin T, Elbin RJ, Harris W, Parker T, Kontos A. *Am J Sports Med*. 2012;40(6):1303-1312.
- Sex and age differences in depression and baseline sport-related concussion neurocognitive performance and symptoms. Covassin T, Elbin RJ 3rd, Larson E, Kontos AP. *Clin J Sport Med*. 2012;22(2):98-104.
- Spectrum of acute clinical characteristics of diagnosed concussions in college athletes wearing instrumented helmets. Duhaime AC, Beckwith JG, Maerlender AC, et al. *J Neurosurg*. 2012.

The utility of post-concussion neuropsychological data in identifying cognitive change following sports-related MTBI in the absence of baseline data. Echemendia RJ, Bruce JM, Bailey CM, Sanders JF, Arnett P, Vargas G. *Clin Neuropsychol*. 2012

Validity of affect measurements in evaluating symptom reporting in athletes. Ramanathan DM, Rabinowitz AR, Barwick FH, Arnett PA. *J Int Neuropsychol Soc*. 2012;18(1):101-107.

## 2011

Assessment and management of sport-related concussions in United States high schools. Meehan WP 3rd, d'Hemecourt P, Collins CL, Comstock RD. *Am J Sports Med*. 2011;39(11):2304-2310.

Assessment of symptoms in a concussion management programme: Method influences outcome. Krol AL, Mrazik M, Naidu D, Brooks BL, Iverson GL. *Brain Inj*. 2011;25(13-14):1300-1305.

Auditory processing after sport-related concussions. Turgeon C, Champoux F, Lepore F, Leclerc S, Ellemberg D. *Ear Hear*. 2011;32(5):667-670.

Baseline neuropsychological testing in managing sport-related concussion: Does it modify risk? Randolph C. *Curr Sports Med Rep*. 2011;10(1):21-26.

Differences in change scores and the predictive validity of three commonly used measures following concussion in the middle school and high school aged population. Barlow M, Schlabach D, Peiffer J, Cook C. *Int J Sports Phys Ther*. 2011;6(3):150-157.

Early indicators of enduring symptoms in high school athletes with multiple previous concussions. Schatz P, Moser RS, Covassin T, Karpf R. *Neurosurgery*. 2011;68(6):1562-7; discussion 1567.

Group versus individual administration affects baseline neurocognitive test performance. Moser RS, Schatz P, Neidzowski K, Ott SD. *Am J Sports Med*. 2011;39(11):2325-2330.

- A history of sport-related concussion on event-related brain potential correlates of cognition. Broglio SP, Moore RD, Hillman CH. *Int J Psychophysiol.* 2011;82(1):16-23.
- Identifying neurocognitive deficits in adolescents following concussion. Thomas DG, Collins MW, Saladino RA, Frank V, Raab J, Zuckerbraun NS. *Acad Emerg Med.* 2011;18(3):246-254
- The immediate post-concussion assessment and cognitive testing battery and traditional neuropsychological measures: A construct and concurrent validity study. Allen BJ, Gfeller JD. *Brain Inj.* 2011;25(2):179-191.
- Long-term neurocognitive dysfunction in sports: What is the evidence? Solomon GS, Ott SD, Lovell MR. *Clin Sports Med.* 2011;30(1):165-77, x-xi.
- Long-term outcomes after uncomplicated mild traumatic brain injury: A comparison with trauma controls. Ponsford J, Cameron P, Fitzgerald M, Grant M, Mikocka-Walus A. *J Neurotrauma.* 2011;28(6):937-946.
- Neuropsychological assessment of sport-related concussion. Johnson EW, Kegel NE, Collins MW. *Clin Sports Med.* 2011;30(1):73-88, viii-ix.
- Neuropsychological testing point/counterpoint. Randolph C, Lovell M, Laker SR. *PM R.* 2011;3(10 Suppl 2):S433-9.
- One-year test-retest reliability of the online version of ImPACT in high school athletes. Elbin RJ, Schatz P, Covassin T. *Am J Sports Med.* 2011;39(11):2319-2324.
- Post-concussion cognitive declines and symptomatology are not related to concussion biomechanics in high school football players. Broglio SP, Eckner JT, Surma T, Kutcher JS. *Post J Neurotrauma.* 2011;28(10):2061-2068.
- Relationship of soccer heading to computerized neurocognitive performance and symptoms among female and male youth soccer players. Kontos AP, Dolese A, Elbin RJ, Covassin T, Warren BL. *Brain Inj.* 2011;25(12):1234-1241.

Sensitivity and specificity of subacute computerized neurocognitive testing and symptom evaluation in predicting outcomes after sports-related concussion. Lau BC, Collins MW, Lovell MR. *Am J Sports Med.* 2011;39(6):1209-1216.

Which on-field signs/symptoms predict protracted recovery from sport-related concussion among high school football players? Lau BC, Kontos AP, Collins MW, Mucha A, Lovell MR. *Am J Sports Med.* 2011;39(11):2311-2318.

## 2010

Case report: Use of the immediate post concussion assessment and cognitive testing (ImPACT) to assist with return to duty determination of special operations soldiers who sustained mild traumatic brain injury. Hettich T, Whitfield E, Kratz K, Frament C. *J Spec Oper Med.* 2010;10(4):48-55.

The cognitive effects and decrements following concussion. Elbin T, Covassin R.J. *Open Access Journal of Sports Medicine.* 2010;1:55-61.

Examination of the construct validity of ImPACT computerized test, traditional, and experimental neuropsychological measures. Maerlender A, Flashman L, Kessler A, et al. *Clin Neuropsychol.* 2010;24(8):1309-1325.

Exploring differences in computerized neurocognitive concussion testing between african american and white athletes. Kontos AP, Elbin RJ 3rd, Covassin T, Larson E. *Arch Clin Neuropsychol.* 2010;25(8):734-744.

Functionally-detected cognitive impairment in high school football players without clinically-diagnosed concussion. Talavage TM, Nauman E, Breedlove EL, et al. *J Neurotrauma.* 2010

High school concussions in the 2008-2009 academic year: Mechanism, symptoms, and management. Meehan WP 3rd, d'Hemecourt P, Comstock RD. *Am J Sports Med.* 2010;38(12):2405-2409.

Influence of poor effort on self-reported symptoms and neurocognitive test performance following mild traumatic brain injury. Lange RT, Iverson GL, Brooks BL, Rennison VL. *J Clin Exp Neuropsychol*. 2010;32(9):961-972.

Investigating baseline neurocognitive performance between male and female athletes with a history of multiple concussion. Covassin T, Elbin R, Kontos A, Larson E. *J Neurol Neurosurg Psychiatry*. 2010;81(6):597-601.

Long-term test-retest reliability of baseline cognitive assessments using ImPACT. Schatz P. *Am J Sports Med*. 2010;38(1):47-53.

Post-concussion symptom reporting and the "good-old-days" bias following mild traumatic brain injury. Lange RT, Iverson GL, Rose A. *Arch Clin Neuropsychol*. 2010;25(5):442-450.

Postconcussive symptoms are associated with compensatory cortical recruitment during a working memory task. Pardini JE, Pardini DA, Becker JT, et al. *Neurosurgery*. 2010;67(4):1020-7; discussion 1027-8.

A prospective study of physician-observed concussions during junior ice hockey: Implications for incidence rates. Echlin PS, Tator CH, Cusimano MD, et al. *Neurosurg Focus*. 2010;29(5):E4.

Relationship between subjective test feedback provided by high-school athletes during computer-based assessment of baseline cognitive functioning and self-reported symptoms. Schatz P, Neidzowski K, Moser RS, Karpf R. *Arch Clin Neuropsychol*. 2010;25(4):285-292.

Return to play after an initial or recurrent concussion in a prospective study of physician-observed junior ice hockey concussions: Implications for return to play after a concussion. Echlin PS, Tator CH, Cusimano MD, et al. *Neurosurg Focus*. 2010;29(5):E5.

Sports-related mild traumatic brain injury in female youths. Keightley ML, Yule A, Garland K, et al. *BMJ Case Rep*. 2010;2010:10.1136/bcr.09.2009.2309.

Supporting the student-athlete's return to the classroom after a sport-related concussion. McGrath N. *J Athl Train*. 2010;45(5):492-498.

Tracking neurocognitive performance following concussion in high school athletes. Covassin T, Elbin RJ, Nakayama Y. *Phys Sportsmed*. 2010;38(4):87-93.

## 2009

The association between mild traumatic brain injury history and cognitive control. Pontifex MB, O'Connor PM, Broglio SP, Hillman CH. *Neuropsychologia*. 2009;47(14):3210-3216.

Computerized neuropsychological profiles of South African versus US athletes: A basis for commentary on cross-cultural norming issues in the sports concussion arena. Shuttleworth-Edwards AB, Whitefield-Alexander VJ, Radloff SE, Taylor AM, Lovell MR. *Phys Sportsmed*. 2009;37(4):45-52.

Concussion in the National Football League: An overview for neurologists. Casson IR, Pellman EJ, Viano DC. *Phys Med Rehabil Clin N Am*. 2009;20(1):195-214, x.

Consensus statement on concussion in sport: The 3rd international conference on concussion in sport held in Zurich, November 2008. McCrory P, Meeuwisse W, Johnston K, et al. *Br J Sports Med*. 2009;43 Suppl 1:i76-90.

Current sport-related concussion teaching and clinical practices of sports medicine professionals. Covassin T, Elbin R 3rd, Stiller-Ostrowski JL. *J Athl Train*. 2009;44(4):400-404.

History of multiple self-reported concussions is not associated with reduced cognitive abilities. Bruce JM, Echemendia RJ. *Neurosurgery*. 2009;64(1):100-6; discussion 106.

Immediate post-concussion assessment and cognitive testing (ImPACT) practices of sports medicine professionals. Covassin T, Elbin RJ 3rd,

- Stiller-Ostrowski JL, Kontos AP. *J Athl Train*. 2009;44(6):639-644.
- The management of sports-related concussion: Current status and future trends. Lovell M. *Clin Sports Med*. 2009;28(1):95-111.
- Neurocognitive and symptom predictors of recovery in high school athletes. Lau B, Lovell MR, Collins MW, Pardini J. *Clin J Sport Med*. 2009;19(3):216-221.
- Neurocognitive evaluation of mild traumatic brain injury in the hospitalized pediatric population. Nance ML, Polk-Williams A, Collins MW, Wiebe DJ. *Ann Surg*. 2009;249(5):859-863.
- Neurocognitive function of emergency department patients with mild traumatic brain injury. Peterson SE, Stull MJ, Collins MW, Wang HE. *Ann Emerg Med*. 2009;53(6):796-803.e1.
- The neurophysiology and assessment of sports-related head injuries. Lovell M. *Phys Med Rehabil Clin N Am*. 2009;20(1):39-53, vii-viii.
- The persistent effects of concussion on neuroelectric indices of attention. Broglio SP, Pontifex MB, O'Connor P, Hillman CH. *J Neurotrauma*. 2009;26(9):1463-1470.
- The relationship of athlete-reported concussion symptoms and objective measures of neurocognitive function and postural control. Broglio SP, Sosnoff JJ, Ferrara MS. *Clin J Sport Med*. 2009;19(5):377-382.
- The role of concussion history and gender in recovery from soccer-related concussion. Colvin AC, Mullen J, Lovell MR, West RV, Collins MW, Groh M. *Am J Sports Med*. 2009;37(9):1699-1704.
- Sports concussion: Management and predictors of outcome. Reddy CC, Collins MW. *Curr Sports Med Rep*. 2009;8(1):10-15.

## 2008

- Biopsychosocial characteristics and neurocognitive test performance in national football league players: An initial assessment. Solomon GS, Haase RF. *Arch Clin Neuropsychol*. 2008;23(5):563-577.



Concussion history and postconcussion neurocognitive performance and symptoms in collegiate athletes. Covassin T, Stearne D, Elbin R. *J Athl Train*. 2008;43(2):119-124.

Concussion in sports: Postconcussive activity levels, symptoms, and neurocognitive performance. Majerske CW, Mihalik JP, Ren D, et al. *J Athl Train*. 2008;43(3):265-274.

Concussion management in the child and adolescent athlete. Lovell MR, Fazio V. *Curr Sports Med Rep*. 2008;7(1):12-15.

Neurocognitive vulnerability amongst university rugby players versus noncontact sport controls. Shuttleworth-Edwards AB, Smith I, Radloff SE. *J Clin Exp Neuropsychol*. 2008;30(8):870-884.

Neuropsychological test performance of Hawaii high school athletes: Hawai'i ImpACT normative data. Tsushima WT, Oshiro R, Zimbra D. *Hawaii Med J*. 2008;67(4):93-95.

## 2007

Comparison of preseason, midseason, and postseason neurocognitive scores in uninjured collegiate football players. Miller JR, Adamson GJ, Pink MM, Sweet JC. *Am J Sports Med*. 2007;35(8):1284-1288.

Differential rate of recovery in athletes after first and second concussion episodes. Slobounov S, Slobounov E, Sebastianelli W, Cao C, Newell K. *Neurosurgery*. 2007;61(2):338-44; discussion 344.

Effectiveness of mouth guards in reducing neurocognitive deficits following sports-related cerebral concussion. Mihalik JP, McCaffrey MA, Rivera EM, et al. *Dent Traumatol*. 2007;23(1):14-20.

Effects of a maximal exercise test on neurocognitive function. Covassin T, Weiss L, Powell J, Womack C. *Br J Sports Med*. 2007;41(6):370-4; discussion 374.

Functional brain abnormalities are related to clinical recovery and time to

- return-to-play in athletes. Lovell MR, Pardini JE, Welling J, et al. *Neurosurgery*. 2007;61(2):352-9; discussion 359-60.
- Hyperbaric oxygen and carbon monoxide poisoning: A critical review. Stoller KP. *Neurol Res*. 2007;29(2):146-155.
- Neurocognitive performance of concussed athletes when symptom free. Broglio SP, Macciocchi SN, Ferrara MS. *J Athl Train*. 2007;42(4):504-508.
- Neuropsychological evaluation in the diagnosis and management of sports-related concussion. Moser RS, Iverson GL, Echemendia RJ, et al. *Arch Clin Neuropsychol*. 2007;22(8):909-916.
- Predicting slow recovery from sport-related concussion: The new simple-complex distinction. Iverson G. *Clin J Sport Med*. 2007;17(1):31-37.
- The relation between post concussion symptoms and neurocognitive performance in concussed athletes. Fazio VC, Lovell MR, Pardini JE, Collins MW. *NeuroRehabilitation*. 2007;22(3):207-216.
- The relationship between neurocognitive function and noncontact anterior cruciate ligament injuries. Swanik CB, Covassin T, Stearne DJ, Schatz P. *Am J Sports Med*. 2007;35(6):943-948.
- Sensitivity of the concussion assessment battery. Broglio SP, Macciocchi SN, Ferrara MS. *Neurosurgery*. 2007;60(6):1050-7; discussion 1057-8.
- Sex differences in neuropsychological function and post-concussion symptoms of concussed collegiate athletes. Covassin T, Schatz P, Swanik CB. *Neurosurgery*. 2007;61(2):345-50; discussion 350-1.
- Test-retest reliability of computerized concussion assessment programs. Broglio SP, Ferrara MS, Macciocchi SN, Baumgartner TA, Elliott R. *J Athl Train*. 2007;42(4):509-514.
- A validation of the post concussion symptom scale in the assessment of complex concussion using cognitive testing and functional MRI. Chen JK, Johnston KM, Collie A, McCrory P, Ptito A. *J Neurol Neurosurg*

*Psychiatry*. 2007;78(11):1231-1238.

2006

The "value added" of neurocognitive testing after sports-related concussion. Van Kampen DA, Lovell MR, Pardini JE, Collins MW, Fu FH. *Am J Sports Med*. 2006;34(10):1630-1635.

Concussion history is not a predictor of computerized neurocognitive performance. Broglio SP, Ferrara MS, Piland SG, Anderson RB, Collie A. *Br J Sports Med*. 2006;40(9):802-5; discussion 802-5.

Concussion in professional football: Helmet testing to assess impact performance--part 11. Pellman EJ, Viano DC, Withnall C, Shewchenko N, Bir CA, Halstead PD. *Neurosurgery*. 2006;58(1):78-96; discussion 78-96.

Concussion in professional football: Recovery of NFL and high school athletes assessed by computerized neuropsychological testing--part 12. Pellman EJ, Lovell MR, Viano DC, Casson IR. *Neurosurgery*. 2006;58(2):263-74; discussion 263-74.

Cross-validation of measures used for computer-based assessment of concussion. Schatz P, Putz BO. *Appl Neuropsychol*. 2006;13(3):151-159.

Die unerkannte hirnerkrankung im sport. Biasca N, Lovell M, Collins M, et al. *Unfallchirurg*. 2006(2):101-111.

Examining concussion rates and return to play in high school football players wearing newer helmet technology: A three-year prospective cohort study. Collins M, Lovell MR, Iverson GL, Ide T, Maroon J. *Neurosurgery*. 2006;58(2):275-86; discussion 275-86.

Measurement of symptoms following sports-related concussion: Reliability and normative data for the post-concussion scale. Lovell MR, Iverson GL, Collins MW, et al. *Appl Neuropsychol*. 2006;13(3):166-174.

MRI and sports concussion: The new window into the brain. Lovell MR, Pardini JE, Collins MW, Fu FH. *Orthopaedic Technology Review*. 2006:24-27.

No cumulative effects for one or two previous concussions. Iverson GL, Brooks BL, Lovell MR, Collins MW. *Br J Sports Med*. 2006;40(1):72-75.

Overview of concussion consensus statements since 2000. Cantu RC, Aubry M, Dvorak J, et al. *Neurosurg Focus*. 2006;21(4):E3.

Recovery from sports concussion in high school and collegiate athletes. McClincy MP, Lovell MR, Pardini J, Collins MW, Spore MK. *Brain Inj*. 2006;20(1):33-39.

Sensitivity and specificity of the ImPACT test battery for concussion in athletes. Schatz P, Pardini JE, Lovell MR, Collins MW, Podell K. *Arch Clin Neuropsychol*. 2006;21(1):91-99.

Tracking neuropsychological recovery following concussion in sport. Iverson GL, Brooks BL, Collins MW, Lovell MR. *Brain Inj*. 2006;20(3):245-252.

Veilig handelen bij sportgerelateerde hersenschuddingen. Van Kampen DA, Lovell MR, Diercks RL. *Geneeskunde en Sport*. 2006;39(4):140-155.

## 2005

Management of cerebral concussion in military personnel: Lessons learned from sports medicine. Lovell M, Collins M, Pardini JE, Parodi A, Yates A. *Oper Tech Sports Med*. 2005;13(4):212-221.

Posttraumatic migraine characteristics in athletes following sports-related concussion. Mihalik JP, Stump JE, Collins MW, Lovell MR, Field M, Maroon JC. *J Neurosurg*. 2005;102(5):850-855.

Prolonged effects of concussion in high school athletes. Moser RS, Schatz P, Jordan BD. *Neurosurgery*. 2005;57(2):300-6; discussion 300-6.

Sex differences in outcome following sports-related concussion. Broshek DK, Kaushik T, Freeman JR, Erlanger D, Webbe F, Barth JT. *J Neurosurg*. 2005;102(5):856-863.

Spinal cord and cerebral concussion in a professional athlete: A case report. Grove R, Lovell MR, Collins MW, Norwig J, Iso A, Maroon JC. *J Athl Train*. 2005;57:56-62.

Standard regression-based methods for measuring recovery after sport-related concussion. McCrea M, Barr WB, Guskiewicz K, et al. *J Int Neuropsychol Soc*. 2005;11(1):58-69.

Summary and agreement statement of the 2nd international conference on concussion in sport, Prague 2004. McCrory P, Johnston K, Meeuwisse W, et al. *Br J Sports Med*. 2005;39(4):196-204.

Validity of ImPACT for measuring processing speed following sports-related concussion. Iverson GL, Lovell MR, Collins MW. *J Clin Exp Neuropsychol*. 2005;27(6):683-689.

## 2004

Athlete concussion management guidelines using functional magnetic resonance imaging. Lovell MR, Collins MW, Fu FH, Stump JE. *Orthopaedic Technology Review*. 2004;6(4).

Concussion history predicts self-reported symptoms before and following a concussive event. Bruce JM, Echemendia RJ. *Neurology*. 2004;63(8):1516-1518.

Concussion in professional football: Neuropsychological testing--part 6. Pellman EJ, Lovell MR, Viano DC, Casson IR, Tucker AM. *Neurosurgery*. 2004;55(6):1290-303; discussion 1303-5.

Cumulative effects of concussion in amateur athletes. Iverson GL, Gaetz M, Lovell MR, Collins MW. *Brain Inj*. 2004;18(5):433-443.

Grade 1 or "ding" concussions in high school athletes. Lovell MR, Collins MW, Iverson GL, Johnston KM, Bradley JP. *Am J Sports Med*. 2004;32(1):47-54.

An introduction to sports concussion for the sport psychologist. Kontos A,

Collins MW, Russo SA. *J Appl Sport Psychol*. 2004;16:220-235.

New developments in the management of sports concussion. Collins MW, Stump JE, Lovell MR. *Curr Opin Orthop*. 2004;15:100-107.

Relation between subjective fogginess and neuropsychological testing following concussion. Iverson GL, Gaetz M, Lovell MR, Collins MW. *J Int Neuropsychol Soc*. 2004;10(6):904-906.

Return to play following sports-related concussion. Lovell M, Collins M, Bradley J. *Clin Sports Med*. 2004;23(3):421-41, ix.

## 2003

Does age play a role in recovery from sports-related concussion? A comparison of high school and collegiate athletes. Field M, Collins MW, Lovell MR, Maroon J. *J Pediatr*. 2003;142(5):546-553.

Epidemiological considerations of concussions among intercollegiate athletes. Covassin T, Swanik CB, Sachs ML. *Appl Neuropsychol*. 2003;10(1):12-22.

Interpreting change on ImPACT following sport concussion. Iverson GL, Lovell MR, Collins MW. *Clin Neuropsychol*. 2003;17(4):460-467.

New technology and sports-related concussion. Lovell MR, Collins MW, Fu FH. *Orthopaedic Technology Review*. 2003;5(1):35-37.

On-field predictors of neuropsychological and symptom deficit following sports-related concussion. Collins MW, Iverson GL, Lovell MR, McKeag DB, Norwig J, Maroon J. *Clin J Sport Med*. 2003;13(4):222-229.

Recovery from mild concussion in high school athletes. Lovell MR, Collins MW, Iverson GL, et al. *J Neurosurg*. 2003;98(2):296-301.

Relationship between postconcussion headache and neuropsychological test performance in high school athletes. Collins MW, Field M, Lovell MR, et al. *Am J Sports Med*. 2003;31(2):168-173.

## 2002

The clinical management of sports concussion. Collins MW, Hawn KL. *Curr Sports Med Rep.* 2002;1(1):12-22.

Cumulative effects of concussion in high school athletes. Collins MW, Lovell MR, Iverson GL, Cantu RC, Maroon JC, Field M. *Neurosurgery.* 2002;51(5):1175-9; discussion 1180-1.

The evaluation of athletes with cerebral concussion. Maroon JC, Field M, Lovell M, Collins M, Bost J. *Clin Neurosurg.* 2002;49:319-332.

New developments in the evaluation of sports-related concussion. Lovell MR, Collins MW. *Curr Sports Med Rep.* 2002;1(5):287-292.

The relevance of neuropsychologic testing for sports-related head injuries. Lovell MR. *Curr Sports Med Rep.* 2002;1(1):7-11.

Summary and agreement statement of the first international conference on concussion in sport, Vienna 2001. Recommendations for the improvement of safety and health of athletes who may suffer concussive injuries. Aubry M, Cantu R, Dvorak J, et al. *Br J Sports Med.* 2002;36(1):6-10.

## 2001

The assessment of sport-related concussion: The evidence behind neuropsychological testing and management. Grindel SH, Lovell MR, Collins MW. *Clin J Sport Med.* 2001;11(3):134-143.

## 1999

Current issues in managing sports-related concussion. Collins MW, Lovell MR, Mckeag DB. *JAMA.* 1999;282(24):2283-2285.

Does loss of consciousness predict neuropsychological decrements after concussion? Lovell MR, Iverson GL, Collins MW, McKeag D, Maroon JC. *Clin J Sport Med.* 1999;9(4):193-198.

Relationship between concussion and neuropsychological performance in college football players. Collins MW, Grindel SH, Lovell MR, et al. *JAMA*. 1999;282(10):964-970.

1998

Neuropsychological assessment of the college football player. Lovell MR, Collins MW. *J Head Trauma Rehabil*. 1998;13(2):9-26.